

***Springer Monographs in Mathematics***

**Springer-Verlag Berlin Heidelberg GmbH**

Jean-Pierre Serre

# Local Algebra

Translated from the French by CheeWhye Chin



Springer

*Jean-Pierre Serre*  
Collège de France  
3 rue d'Ulm  
75005 Paris, France  
e-mail: serre@dmi.ens.fr

*Translator:*  
CheeWhye Chin  
Princeton University  
Department of Mathematics  
Princeton, NJ 08544  
USA  
e-mail: cchin@princeton.edu

CIP data applied for

Die Deutsche Bibliothek - CIP-Einheitsaufnahme

**Serre, Jean-Pierre:**  
Local algebra / Jean-Pierre Serre. Transl. from the French by CheeWhye Chin. - Berlin; Heidelberg;  
New York; Barcelona; Hong Kong; London; Milan; Paris; Singapore; Tokyo: Springer, 2000  
(Springer monographs in mathematics)  
Einheitssacht.: Algèbre locale <engl.>  
ISBN 978-3-642-08590-1      ISBN 978-3-662-04203-8 (eBook)  
DOI 10.1007/978-3-662-04203-8

---

Mathematics Subject Classification (2000): 13xx

---

ISSN 1439-7382  
ISBN 978-3-642-08590-1

This work is subject to copyright. All rights are reserved, whether the whole or part of the material is concerned, specifically the rights of translation, reprinting, reuse of illustrations, recitation, broadcasting, reproduction on microfilm or in any other way, and storage in data banks. Duplication of this publication or parts thereof is permitted only under the provisions of the German Copyright Law of September 9, 1965, in its current version, and permission for use must always be obtained from Springer-Verlag. Violations are liable for prosecution under the German Copyright Law.

© Springer-Verlag Berlin Heidelberg 2000  
Originally published by Springer-Verlag Berlin Heidelberg New York in 2000  
Softcover reprint of the hardcover 1st edition 2000

The use of general descriptive names, registered names, trademarks etc. in this publication does not imply, even in the absence of a specific statement, that such names are exempt from the relevant protective laws and regulations and therefore free for general use.

Cover design: *Erich Kirchner, Heidelberg*  
Typesetting by the translator using a Springer  $\LaTeX$  macro package  
Printed on acid-free paper SPIN 10703977      41/3143ck-5 4 3 2 1 0

# Preface

The present book is an English translation of

*Algèbre Locale — Multiplicités*

published by Springer-Verlag as no. 11 of the Lecture Notes series.

The original text was based on a set of lectures, given at the Collège de France in 1957-1958, and written up by Pierre Gabriel. Its aim was to give a short account of Commutative Algebra, with emphasis on the following topics:

- a) *Modules* (as opposed to *Rings*, which were thought to be the only subject of Commutative Algebra, before the emergence of sheaf theory in the 1950s);
- b) *Homological methods*, à la Cartan-Eilenberg;
- c) *Intersection multiplicities*, viewed as Euler-Poincaré characteristics.

The English translation, done with great care by CheeWhye Chin, differs from the original in the following aspects:

- The terminology has been brought up to date (e.g. “cohomological dimension” has been replaced by the now customary “depth”).
- I have rewritten a few proofs and clarified (or so I hope) a few more.
- A section on graded algebras has been added (App. III to Chap. IV).
- New references have been given, especially to other books on Commutative Algebra: Bourbaki (whose Chap. X has now appeared, after a 40-year wait), Eisenbud, Matsumura, Roberts, . . . .

I hope that these changes will make the text easier to read, without changing its informal “Lecture Notes” character.

J-P. Serre,  
Princeton, Fall 1999

# Contents

<b>Preface</b> .....	<b>v</b>
<b>Contents</b> .....	<b>vii</b>
<b>Introduction</b> .....	<b>xi</b>
<b>I. Prime Ideals and Localization</b> .....	<b>1</b>
§1. Notation and definitions .....	1
§2. Nakayama's lemma .....	1
§3. Localization .....	2
§4. Noetherian rings and modules .....	4
§5. Spectrum .....	4
§6. The noetherian case .....	5
§7. Associated prime ideals .....	6
§8. Primary decompositions .....	10
<b>II. Tools</b> .....	<b>11</b>
A: Filtrations and Gradings .....	11
§1. Filtered rings and modules .....	11
§2. Topology defined by a filtration .....	12
§3. Completion of filtered modules .....	13
§4. Graded rings and modules .....	14
§5. Where everything becomes noetherian again — $q$ -adic filtrations .....	17
B: Hilbert-Samuel Polynomials .....	19
§1. Review on integer-valued polynomials .....	19
§2. Polynomial-like functions .....	21
§3. The Hilbert polynomial .....	21
§4. The Samuel polynomial .....	24

<b>III. Dimension Theory</b> .....	<b>29</b>
A: Dimension of Integral Extensions .....	29
§1. Definitions .....	29
§2. Cohen-Seidenberg first theorem .....	30
§3. Cohen-Seidenberg second theorem .....	32
B: Dimension in Noetherian Rings .....	33
§1. Dimension of a module .....	33
§2. The case of noetherian local rings .....	33
§3. Systems of parameters .....	36
C: Normal Rings .....	37
§1. Characterization of normal rings .....	37
§2. Properties of normal rings .....	38
§3. Integral closure .....	40
D: Polynomial Rings .....	40
§1. Dimension of the ring $A[X_1, \dots, X_n]$ .....	40
§2. The normalization lemma .....	42
§3. Applications. I. Dimension in polynomial algebras ....	44
§4. Applications. II. Integral closure of a finitely generated algebra .....	46
§5. Applications. III. Dimension of an intersection in affine space .....	47
<b>IV. Homological Dimension and Depth</b> .....	<b>51</b>
A: The Koszul Complex .....	51
§1. The simple case .....	51
§2. Acyclicity and functorial properties of the Koszul complex .....	53
§3. Filtration of a Koszul complex .....	56
§4. The depth of a module over a noetherian local ring ...	59
B: Cohen-Macaulay Modules .....	62
§1. Definition of Cohen-Macaulay modules .....	63
§2. Several characterizations of Cohen-Macaulay modules .....	64
§3. The support of a Cohen-Macaulay module .....	66
§4. Prime ideals and completion .....	68
C: Homological Dimension and Noetherian Modules .....	70
§1. The homological dimension of a module .....	70
§2. The noetherian case .....	71
§3. The local case .....	73

D: Regular Rings .....	75
§1. Properties and characterizations of regular local rings .....	75
§2. Permanence properties of regular local rings .....	78
§3. Delocalization .....	80
§4. A criterion for normality .....	82
§5. Regularity in ring extensions .....	83
Appendix I: Minimal Resolutions .....	84
§1. Definition of minimal resolutions .....	84
§2. Application .....	85
§3. The case of the Koszul complex .....	86
Appendix II: Positivity of Higher Euler-Poincaré Characteristics .....	88
Appendix III: Graded-polynomial Algebras .....	91
§1. Notation .....	91
§2. Graded-polynomial algebras .....	92
§3. A characterization of graded-polynomial algebras .....	93
§4. Ring extensions .....	93
§5. Application: the Shephard-Todd theorem .....	95
<b>V. Multiplicities .....</b>	<b>99</b>
A: Multiplicity of a Module .....	99
§1. The group of cycles of a ring .....	99
§2. Multiplicity of a module .....	100
B: Intersection Multiplicity of Two Modules .....	101
§1. Reduction to the diagonal .....	101
§2. Completed tensor products .....	102
§3. Regular rings of equal characteristic .....	106
§4. Conjectures .....	107
§5. Regular rings of unequal characteristic (unramified case) .....	108
§6. Arbitrary regular rings .....	110
C: Connection with Algebraic Geometry .....	112
§1. Tor-formula .....	112
§2. Cycles on a non-singular affine variety .....	113
§3. Basic formulae .....	114
§4. Proof of theorem 1 .....	116
§5. Rationality of intersections .....	116



§6. Direct images ..... 117  
§7. Pull-backs ..... 117  
§8. Extensions of intersection theory ..... 119

**Bibliography ..... 123**

**Index ..... 127**

**Index of Notation ..... 129**

# Introduction

The intersection multiplicities of algebraic geometry are equal to some “Euler-Poincaré characteristics” constructed by means of the Tor functor of Cartan-Eilenberg. The main purpose of this course is to prove this result, and to apply it to the fundamental formulae of intersection theory.

It is necessary to first recall some basic results of local algebra: primary decomposition, Cohen-Seidenberg theorems, normalization of polynomial rings, Krull dimension, characteristic polynomials (in the sense of Hilbert-Samuel).

Homology comes next, when we consider the multiplicity  $e_{\mathfrak{q}}(E, r)$  of an ideal of definition  $\mathfrak{q} = (x_1, \dots, x_r)$  of a local noetherian ring  $A$  with respect to a finitely generated  $A$ -module  $E$ . This multiplicity is defined as the coefficient of  $n^r/r!$  in the polynomial-like function  $n \mapsto \ell_A(E/\mathfrak{q}^n E)$  [here  $\ell_A(F)$  is the length of an  $A$ -module  $F$ ]. We prove in this case the following formula, which plays an essential role in the sequel:

$$e_{\mathfrak{q}}(E, r) = \sum_{i=0}^r (-1)^i \ell_A(H_i(\mathbf{x}, E)) \quad (*)$$

where the  $H_i(\mathbf{x}, E)$  denotes the homology modules of the Koszul complex constructed on  $E$  by means of  $\mathbf{x} = (x_1, \dots, x_r)$ .

Moreover this complex can be used in other problems of local algebra, for example for the study of the depth of modules over a local ring and of the Cohen-Macaulay modules (those whose Krull dimension coincides with their depth), and also for showing that regular local rings are the only local rings whose homological dimension is finite.

Once formula (\*) is proved, one may study the Euler-Poincaré characteristic constructed by means of Tor. When one translates the geometric situation of intersections into the language of local algebra, one obtains a regular local ring  $A$ , of dimension  $n$ , and two finitely generated  $A$ -modules  $E$  and  $F$  over  $A$ , whose tensor product is of finite length over  $A$  (this means that the varieties corresponding to  $E$  and  $F$  intersect only at the given point). One is then led to conjecture the following statements:

- (i)  $\dim(E) + \dim(F) \leq n$  (“dimension formula”).
- (ii)  $\chi_A(E, F) = \sum_{i=0}^n (-1)^i \ell_A(\text{Tor}_i^A(E, F))$  is  $\geq 0$ .
- (iii)  $\chi_A(E, F) = 0$  if and only if the inequality in (i) is strict.

Formula (\*) shows that the statements (i), (ii) and (iii) are true if  $F = A/(x_1, \dots, x_r)$ , with  $\dim(F) = n - r$ . Thanks to a process, using completed tensor products, which is the algebraic analogue of “reduction to the diagonal”, one can show that they are true when  $A$  has the same characteristic as its residue field, or when  $A$  is unramified. To go beyond that, one can use the structure theorems of complete local rings to prove (i) in the most general case. On the other hand, I have not succeeded in proving (ii) and (iii) without making assumptions about  $A$ , nor to give counter-examples. It seems that it is necessary to approach the question from a different angle, for example by directly defining (by a suitable asymptotic process) an integer  $\geq 0$  which one would subsequently show to be equal to  $\chi_A(E, F)$ .

Fortunately, the case of equal characteristic is sufficient for the applications to algebraic geometry (and also to analytic geometry). More specifically, let  $X$  be a non-singular variety, let  $V$  and  $W$  be two irreducible subvarieties of  $X$ , and suppose that  $C = V \cap W$  is an irreducible subvariety of  $X$ , with:

$$\dim X + \dim C = \dim V + \dim W \quad (\text{“proper” intersection}).$$

Let  $A, A_V, A_W$  be the local rings of  $X, V$  and  $W$  at  $C$ . If

$$i(V \cdot W, C; X)$$

denotes the multiplicity of the intersection of  $V$  and  $W$  at  $C$  (in the sense of Weil, Chevalley, Samuel), we have the formula:

$$i(V \cdot W, C; X) = \chi_A(A_V, A_W). \quad (**)$$

This formula is proved by reduction to the diagonal, and the use of (\*). In fact, it is convenient to take (\*\*) as the definition of multiplicities. The properties of these multiplicities are then obtained in a natural way: commutativity follows from that for  $\text{Tor}$ ; associativity follows from the two spectral sequences which expresses the associativity of  $\text{Tor}$ ; the projection formula follows from the two spectral sequences connecting the direct images of a coherent sheave and  $\text{Tor}$  (these latter spectral sequences have other interesting applications, but they are not explored in the present course). In each case, one uses the well-known fact that Euler-Poincaré characteristics remain constant through a spectral sequence.

When one defines intersection multiplicities by means of the  $\text{Tor}$ -formula above, one is led to extend the theory beyond the strictly “non-singular” framework of Weil and Chevalley. For example, if  $f : X \rightarrow Y$

is a morphism of a variety  $X$  into a non-singular variety  $Y$ , one can associate, to two cycles  $x$  and  $y$  of  $X$  and  $Y$ , a “product”  $x \cdot_f y$  which corresponds to  $x \cap f^{-1}(y)$  (of course, this product is only defined under certain dimension conditions). When  $f$  is the identity map, one recovers the standard product. The commutativity, associativity and projection formulae can be stated and proved for this new product.